DISCLAIMER

This paper was submitted to the *Memorias do Instituto Oswaldo Cruz* on 24 January 2018 and was posted to the Fast Track site on 31 January 2018. The information herein

is available for unrestricted use, distribution and reproduction provided that the original work is properly cited as indicated by the Creative Commons Attribution licence (CC

BY).

RECOMMENDED CITATION

de Mendonça MCL, Mares-Guia MA, Rodrigues CDS, dos Santos CC, Chalhoub FLL, Araújo ESM, et al. Imported case of dengue virus 3 genotype I in Rio de Janeiro state,

Brazil [Submitted]. Mem Inst Oswaldo Cruz E-pub: 31 Jan 2018. doi: 10.1590/0074-

02760180036.

Running title: DENV-3 Genotype I in Rio de Janeiro

Title: Imported case of Dengue virus 3 Genotype I in Rio de Janeiro state, Brazil

Author's names:

Marcos César Lima de Mendonça¹, Maria Angelica Mares-Guia¹, Cintia Damasceno

dos Santos Rodrigues¹, Carolina Cardoso dos Santos¹, Flavia Lowen Levy Chalhoub¹, Eliane Saraiva Machado Araújo¹, Alexandre Otávio Chieppe², Rita Maria Ribeiro

Nogueira¹, Ana Maria Bispo de Filippis¹

Institutional affiliations:

¹ Fundação Oswaldo Cruz-Fiocruz, Instituto Oswaldo Cruz, Laboratório de Flavivírus,

Rio de Janeiro, RJ, Brazil.

² Secretaria Estadual de Saúde do Rio de Janeiro

Corresponding author:

Marcos César Lima de Mendonça, marcosclm@ioc.fiorcuz.br

Abstracts:

The dengue virus (DENV), of genus Flavivirus (Flaviviridae), has four antigenically

distinct serotypes, of which DENV-3 is classified into five genotypes. Here, we describe

the detection of DENV-3 genotype I in sera of a Brazilian patient returning from

Singapore to Rio de Janeiro, Brazil by using multiplex real-time RT-PCR, Sanger

sequencing of the whole envelope protein, and phylogenetic analysis. The virus shares

ancestry with those identified in Bali, Indonesia in 2015. Because of tourism and mass

events occurring in the last four years (World Youth Day, World Cup, Olympics),

arboviruses, such as Chikungunya genotype ECSA, DENV-4 Genotype I, and Zika,

from other continents have been introduced in Brazil.

Key words: Dengue, Sequencing, Genotyping

Sponsorships: CNPq, FAPERJ

Dengue virus (DENV) is considered to be the most important arbovirus worldwide and approximately two-fifths of the world's population live in areas endemic for dengue (Guo et al. 2017). DENV is classified into family *Flaviviridae*, genus *Flavivirus*, and comprises four antigenically distinct serotype, Dengue virus 1 (DENV-1), Dengue virus 2 (DENV-2), Dengue virus 3 (DENV-3), and Dengue virus 4 (DENV-4), transmitted to humans by mosquitoes belonging to genus *Aedes*, mainly *Aedes aegypti*, prevalent in tropical and subtropical areas (Lanciotti et al. 1994, Whitehead RH et al. 1971).

Phylogenetic studies focusing the envelope protein proposed the further classification of four genotypes of DENV-3 (genotypes I, II, III, and IV). The first discovered genotype of DENV-3, Genotype IV, identified in Caribbean and Puerto Rico in 1963 (Lanciotti et al. 1994), was the only one circulating in different countries in the Americas, but may be currently extinct. Viruses classified as Genotype I were found in Colombia in 2003 (Usme-Ciro et al. 2008). In Brazil, similar viruses were detected in 2003, but the authors proposed a new genotype (genotype V) for the correct classification of these viruses owing to their topology in the phylogenetic tree. According to this classification, Genotype V viruses were present in Brazil and Colombia between 2003 and 2006. They were closely related to viruses from Asian samples isolated in around 1956, and until that date there was no record of their presence in nature (Aquino et al. 2009). Genotype III was introduced in Latin America in around 1991 (Araújo et al. 2009) and in Brazil in 2001 (Miagostovich et al. 2002), becoming the predominant genotype till date. To our knowledge, there is no record of the circulation of genotype II in the Americas.

Out of the 733 acute sera tested for dengue by real time RT-PCR using the protocol described by Santiago et al., only one sample was positive for DENV-3

presenting a Ct value of 23. The sample was obtained from a 36-year-old Brazilian, who lives in Singapore and toured Indonesia and other Asian countries during a 30-day vacation. On July 3rd, 2017, the patient arrived in Brazil with symptoms of dengue. The patient sought medical assistance and stayed at home, in an urban area of Rio de Janeiro state, for two weeks until the symptoms disappeared.

The Qiagen Onestep RT-PCR Kit was used to generate a 1795-bp fragment, which contains the entire coding region for the envelope protein. Ten primers were used for Sanger sequencing reactions (Data not shown; Acknowledgements). The MEGA 7 software package was employed for choosing the best model of nucleotide substitution and construction of the phylogenetic trees.

The tree shown in figure A was constructed using 2,040 sequences, obtained by Blast, of the envelope protein coding region from all five genotypes. In figure A, the virus from the municipality of residence of the patient, identified as Brazil-RJ/2017/LABFLA (MG812331), belongs to genotype I among samples from Indonesia and Singapore. It is in a cluster supported by 86 of bootstrap with four other samples, which are described in a study with samples from Bali, Indonesia, collected in 2015 (Megawati et al. 2017) Therefore, viruses from Bali, Indonesia and those from the patient (Brasil-RJ/2017/LABFLA) in this study have common ancestry.

A second tree (B) was constructed because there were no complete envelope protein sequence deposits from the Colombian samples classified as genotype I, reclassified as genotype V. The second tree was assembled with 224 3'-terminal nucleotides of the envelope protein coding region. It is evident by the phylogenetic tree that its introduction in Colombia has no correlation with the virus sample found in Rio de Janeiro. The patient probably acquired the DENV-3 infection during his stay in

Indonesia, arriving to Brazil infected. The surveillance in the area around the patient's residence during the viremic phase is very important, to identify other probable cases.

Again, our results show that Brazil has become an important gateway to arboviruses from other continents, as seen with Zika, Yellow Fever, Chikungunya genotype ECSA, DENV-4 Genotype I, and maybe DENV-3 Genotype I recently. This may be hiding other arboviruses or new genotypes. The identification of DENV-3 genotype I in this patient is an indication for the Brazilian health authorities to verify possible changes in DENV-3 infection pattern.

Acknowledgements

This study was partially supported by the ViGenDA's Project, Vigilancia Genómica del dengue, (Genomics Dengue Surveillance), from PAHO/WHO Dengue Regional Program, PAHO RELDA (PAHO Arbovirus Laboratory Network) and its partner, CDC Dengue branch, Puerto Rico, who provided the protocol and primers for sequencing used in this study.

Authors' contribution

Marcos César Lima de Mendonça, carried out DNA Sequencing, phylogenetic analyses, and wrote drafted the manuscript.

Maria Angelica Mares-Guia, Cintia Damasceno dos Santos Rodrigues, Carolina Cardoso dos Santos, Flavia Lowen Levy Chalhoub, Eliane Saraiva Machado Araújo, Alexandre Otávio Chieppe, carried out the laboratory experiments.

Rita Maria Ribeiro Nogueira, Ana Maria Bispo de Filippis designed the study.

References

Aquino VH, Amarilla AA, Alfonso HL, Batista WC, Figueiredo LT. New genotype of dengue type 3 virus circulating in Brazil and Colombia showed a close relationship to old Asian viruses. PLoS One. 2009; 4(10): e7299.

Araújo JM1, Nogueira RM, Schatzmayr HG, Zanotto PM, Bello G. Phylogeography and evolutionary history of dengue vírus type 3. Infect Genet Evol. 2009; 9(4): 716-25.

Guo C, Zhou Z, Wen Z, Liu Y, Zeng C, Xiao D, Ou M, Han Y, Huang S, Liu D, Ye X, Zou X, Wu J, Wang H, Zeng EY, Jing C, Yang G. Global epidemiology of dengue outbreaks in 1990-2015: a systematic review and meta-Analysis. Front Cell Infect Microbiol. 2017; 7: 317.

Lanciotti RS, Lewis JG, Gubler DJ, Trent DW. Molecular evolution and epidemiology of dengue-3 viruses. J Gen Virol. 1994; 75: 65-75.

Miagostovich MP1, dos Santos FB, de Simone TS, Costa EV, Filippis AM, Schatzmayr HG, Nogueira RM. Genetic characterization of dengue virus type 3 isolates in the State of Rio de Janeiro, 2001. Braz J Med Biol Res. 2002; 35(8): 869-72.

Megawati D, Masyeni S, Yohan B, Lestarini A, Hayati RF, Meutiawati F, Suryana K, Widarsa T, Budiyasa DG, Budiyasa N, Myint KSA, Sasmono RT. Dengue in Bali:

Clinical characteristics and genetic diversity of circulating dengue viruses. PLoS Negl Trop Dis. 2017; 11(5): e0005483.

Santiago GA, Vergne E, Quiles Y, Cosme J, Vazquez J, Medina JF, Medina F, Colón C, Margolis H, Muñoz-Jordán JL. Analytical and clinical performance of the CDC real time RT-PCR assay for detection and typing of dengue virus. PLoS Negl Trop Dis. 2013; 7(7): e2311.

Usme-Ciro JA, Mendez JA, Tenorio A, Rey GJ, Domingo C, Gallego-Gomez JC. Simultaneous circulation of genotypes I and III of dengue virus 3 in Colombia. Virol J. 2008; 5: 101.

Whitehead RH, Yuill TM, Gould DJ, Simasathien P. Experimental infection of *Aedes aegypti* and *Aedes albopictus* with dengue viruses. Trans R Soc Trop Med Hyg. 1971; 65(5): 661-7.




Figure 1

Phylogenetic analyses based on complete (A) and partial (B – 224-bp 3- terminal) Protein E coding region sequences were performed in MEGA 7. Closed black circle represents the strain identified in the patient (Brasil-RJ/2017/LABFLA). The trees were inferred with 1000 bootstrap replicates, using maximum likelihood algorithm based on GTR model with Gamma distribution and invariant site correction.